

EVERCYTE GmbH the pharmacocellomics<sup>™</sup> company

Muthgasse 18 1190 Wien

.....

office@evercyte.com www.evercyte.com

# Protocol for cryopreservation of NHEK/SVTERT3-5 cells

Version: April 2017

## Reagents

Trypsin-EDTA (Gibco, Cat# 25300054) Trypsin-Inhibitor (Gibco, Cat# R007100) Freezing medium which contains:

- KGM<sup>™</sup>-2 BulletKit<sup>™</sup> without GA-1000 from this kit
- 10 % DMSO

Storage temperature: liquid nitrogen

- KeratinoUp ready-to-use medium (Evercyte, Cat# MHT-011) which contains:
  - KGM<sup>™</sup>-2 BulletKit<sup>™</sup> (Lonza, Cat# CC-3107) without GA-1000 from this kit and
  - 50 µg/ml G418 (InvivoGen, Cat# ant-gn-5)

Ethanol, 70 %

Practical application

## Freezing of cells:

Detach the cells from culture vessel by using Trypsin-EDTA and Trypsin-Inhibitor as described in Evercyte's protocol for in vitro propagation of NHEK/SVTERT3-5 cells, resuspend detached cells in growth medium and centrifuge at 170 g for 5 min. Then, discard the supernatant, resuspend the cell pellet in the remaining droplet and add freezing medium (4°C) to reach a cell density of about  $5 \times 10^5$  cells/ml (for thawing in a 25 cm<sup>2</sup> culture flask). Add 1 ml of this cell suspension to each pre-cooled cryovial and immediately transfer the cells to -80°C. After 24 hours transfer the vials to liquid nitrogen.

## Thawing of cells:

Add 6 ml of cultivation medium to a 25 cm<sup>2</sup> culture flask and place the culture flask in the incubator for at least 30 min to allow the medium to reach its normal pH. Take a vial of frozen cells, rinse outside with ethanol and pre-warm in hand until one last piece of frozen cells is seen. Then, immediately transfer the content of the vial to a 15 ml centrifugation tube pre-filled with 9 ml of medium pre-cooled to 4°C and centrifuge for 5 min at 170 g. Then, discard the supernatant and resuspend the cells in the remaining droplet. Add 1 ml of pre-warmed medium to the cells, transfer them to the prepared culture flask and incubate at 37°C in a suitable incubator.

Perform a medium change 24 hours after thawing. If the cells are already confluent at this point, they should be passaged (see Evercyte's protocol for in vitro propagation of NHEK/SVTERT3-5 cells).

## Related products

• KeratinoUp ready-to-use medium, 500 ml (Evercyte, Cat# MHT-011)